

第1表 日本の上部地殻の平均組成 (Togashi et al., 2000)

										元素成分率	
SiO ₂ (%)	67.53	Sc (ppm)	16	Y (ppm)	26	Sm (ppm)	4.28	Hf (ppm)	4.1	O	48.06
TiO ₂ (%)	0.62	V (ppm)	110	Zr (ppm)	135	Eu (ppm)	1.07	Ta (ppm)	0.72	Si	31.56
Al ₂ O ₃ (%)	14.67	Cr (ppm)	84	Nb (ppm)	9	Gd (ppm)	3.65	Pb (ppm)	16.9	Ti	0.37
Fe ₂ O ₃ (%)	5.39	Co (ppm)	15	Sb (ppm)	0.61	Tb (ppm)	0.7	Th (ppm)	8.3	Al	7.76
MnO (%)	0.11	Ni (ppm)	38	Cs (ppm)	5.5	Dy (ppm)	3.94	U (ppm)	2.32	Fe	3.77
MgO (%)	2.53	Cu (ppm)	25	Ba (ppm)	458	Ho (ppm)	0.72			Mn	0.09
CaO (%)	3.9	Zn (ppm)	74.1	La (ppm)	21.7	Er (ppm)	2.39			Mg	1.53
Na ₂ O (%)	2.72	As (ppm)	6.5-7.1	Ce (ppm)	46.4	Tm (ppm)	0.39			Ca	2.79
K ₂ O (%)	2.42	Rb (ppm)	85	Pr (ppm)	5.54	Yb (ppm)	2.57			Na	2.02
P ₂ O ₅ (%)	0.12	Sr (ppm)	225	Nd (ppm)	20.8	Lu (ppm)	0.4			K	2.01
										P	0.05